## Specific Mixing Rules: Test and Application

A. Müller, D. Heene-Medina, M. Wendland, J. Vrabec, J. Winkelmann, and J. Fischer

<sup>1</sup> Institut für Physikalische Chemie Universität Halle-Wittenberg D-06217 Merseburg, Germany

<sup>2</sup> Institut für Thermo- und Fluiddynamik Ruhr-Universität Bochum D-44780 Bochum, Germany

<sup>3</sup> Institut für Land-, Umwelt-, und Energietechnik Universität für Bodenkultur A-1190 Wien, Austria

We start from the Helmholtz energy F being written as the sum  $F = F_H + F_A + F_D$  where  $F_H$  is the hard-body contribution,  $F_A$  the attractive dispersion force contribution and  $F_D$  the dipolar contribution. In each of these terms some substance specific parameters like a characteristic density or a characteristic temperature occur. In the case of mixtures we need mixing rules for these parameters. If different sets of mixing rules are used in  $F_H$ ,  $F_A$  and  $F_D$  we call them specific mixing rules." A specific mixing rule for  $F_H$  was given by Boublik and Mansoori. Here, we discuss specific mixing rules for  $F_A$  and  $F_D$ . For  $F_A$ , several specific mixing rules will be tested for Lennard-Jones model mixtures by comparison with molecular simulation results. In particular, these tests are performed for vapor-liquid-phase-equilibria obtained with the NpT + test particle method. For  $F_D$ , we show again by comparison with simulation data that one mixing rule derived from perturbation theory works very well. Using these validated mixing rules, we enter the recently presented BACKONE family of equations of state and present thermodynamic properties including phase equilibria for the two binary mixtures methane + ethane and xenon + methylflouride. It is remarkable that only one state-independent parameter is needed for the description of each mixture.